

Determination of Pesticides in Red Wine by QuEChERS Extraction, Quick QuEChERS Clean-up, and LC/MS/MS Detection

UCT Part Numbers:

RFV0050CT (50 mL polypropylene centrifuge tube) **ECQUUS2-MP (**Mylar Pouch contains: 4000 mg MgSO₄, 2000 mg NaCl) **ECPURMPSMC** (Quick QuEChERS cartridge, 110 mg MgSO₄, 180 mg PSA)

The analysis of pesticide residues in red wines is challenging due to the complexity of the matrix, which contains organic acids, sugars, phenols, and pigments, such as anthocyanins. A simple, faster, and easy to use method is developed for the determination of pesticide residues in red wines.

Eight pesticides with a wide range of polarities (LogP from -0.779 to 5.004) were selected as target analytes. Excellent accuracy and precision data were achieved using this method. Recoveries of planar pesticides, such as Carbendazim and Thiabendazole were not affected since PSA was used for clean-up instead of GCB. PSA removed organic acids, sugars and pigments from the red wine extract. Six red wine samples were extracted using this method. Cyprodinil and Carbendazim were detected in the red wine samples tested, with minimum reporting limits of 1.5 ng/mL.

Procedure

1. Extraction

- a) Add 10 mL of red wine sample to a 50 mL polypropylene centrifuge tube (**RFV0050CT**)
- b) Spike with the appropriate amount of target analytes for fortified samples
- c) Vortex 30 sec, then equilibrate for 15 min
- d) Add 10 mL of acetonitrile, vortex 30 sec
- e) Add salts in Mylar pouch (ECQUUS2-MP)
- f) Shake vigorously for 1 min
- g) Centrifuge at 5000 rpm for 5 min at 20° C
- h) Supernatant is ready for clean-up

2. Quick QuEChERS Clean-up

- a) Draw 1 mL of supernatant into a disposable polypropylene syringe
- b) Pass the supernatant slowly through the Quick QuEChERS cartridge (ECPURMPSMC)

- c) Collect 0.5 mL of the cleaned extract into a 2 mL auto-sampler vial
- d) Add 10 µL 5 ppm TPP as internal standard (IS)
- e) Samples are ready for LC/MS/MS analysis

Clean-up red wine extract with Quick QuEChERS



Quick QuEChERS before (left) and after (right) clean-up of 1 mL red wine extract



3. LC/MS/MS Detection

LC: Thermo Accela 1250 pump with PAL auto-sampler

| LC Conditions | | | | | | | |
|--------------------|---|--|--|--|--|--|--|
| Column | Guard column: Restek C18, 2.1 x 20 mm Column: Sepax HP-C18, 2.1 x 100 mm, 3 µm, 120 Å | | | | | | |
| Column Temperature | Ambient | | | | | | |
| Injection Volume | 10 μL at 15° C | | | | | | |
| Mobile Phase | A: 0.1% formic acid in Milli-Q-water B: 0.1% formic acid in methanol | | | | | | |
| Flow Rate | 200 µL/min | | | | | | |

| Time | % A | %В |
|------|------------|----|
| 0 | 95 | 5 |
| 1 | 95 | 5 |
| 3 | 50 | 50 |
| 8 | 5 | 95 |
| 14.2 | 95 | 5 |
| 16 | 95 | 5 |

-1:

MS/MS: Thermo TSQ Vantage tandem MS

| MS Conditions | | | | | | |
|-------------------------------------|-------------------------|--|--|--|--|--|
| Ion source: | Heated ESI | | | | | |
| Ion polarity: | ESI + | | | | | |
| Spray voltage: | 3000 V | | | | | |
| Sheath gas pressure: | N ₂ @ 40 psi | | | | | |
| Auxiliary gas pressure: | N ₂ @ 10 psi | | | | | |
| Ion transfer capillary temperature: | 350 °C | | | | | |
| Scan type: | SRM (0-16 min) | | | | | |
| CID conditions: | Ar @ 1.5 mTorr | | | | | |

| CDM | 4 | 11.000 |
|--------------|-------|--------|
| SKIVI | trans | itions |
| ••••• | | |

| Compound | Parent | Product ion 1 | CE | Product ion 2 | CE | S-Lens | Dwell time (s) |
|---------------|---------|------------------|----|------------------|----|--------|-------------------|
| Methamidophos | 142.044 | 94.090 | 14 | 125.050 | 16 | 59 | 0.15 |
| Carbendazim | 192.093 | 132.080 | 29 | 160.080 | 17 | 81 | 0.10 |
| Thiabendazole | 202.059 | 131.060 | 31 | 175.070 | 31 | 103 | 0.10 |
| Pyrimethanil | 200.116 | 107.060 | 23 | 183.140 | 22 | 66 | 0.10 |
| Cyprodinil | 226.122 | 77.030 | 40 | 93.050 | 33 | 88 | 0.10 |

| TPP (IS) | 327.093 | 77.020 | 37 | 152.070 | 33 | 98 | 0.10 |
|--------------|---------|---------|----|---------|----|-----|------|
| Diazinon | 305.135 | 153.090 | 15 | 169.08 | 14 | 89 | 0.10 |
| Pyrazophos | 374.103 | 194.060 | 20 | 222.130 | 20 | 104 | 0.10 |
| Chlorpyrifos | 349.989 | 96.890 | 32 | 197.940 | 17 | 69 | 0.10 |

Matrix matched calibration, LOD and LOQ

| Compound | Linearity range (ng/mL) | R ² | LOD (ng/mL) | LOQ (ng/mL) |
|---------------|----------------------------|----------------|----------------|----------------|
| Methamidophos | 2-400 | 0.9991 | 0.15 | 0.49 |
| Carbendazim | 2-400 | 0.9981 | 0.40 | 1.33 |
| Thiabendazole | 2-400 | 0.9940 | 0.09 | 0.31 |
| Pyrimethanil | 2-400 | 0.9990 | 0.01 | 0.05 |
| Cyprodinil | 2-400 | 0.9995 | 0.17 | 0.57 |
| Diazinon | 2-400 | 0.9982 | 0.06 | 0.21 |
| Pyrazophos | 2-400 | 0.9976 | 0.08 | 0.27 |
| Chlorpyrifos | 2-400 | 0.9981 | 0.10 | 0.32 |

Accuracy and Precision Data

| Compound | Fortified at 10 ng/mL | | Fortified at 5 | 0 ng/mL | Fortified at 100 ng/mL | |
|---------------|-----------------------|------------|----------------|------------|------------------------|------------|
| | Recovery% | RSD% (n=4) | Recovery% | RSD% (n=4) | Recovery% | RSD% (n=4) |
| Methamidophos | 93.7 | 3.4 | 81.6 | 5.8 | 84.2 | 3.5 |
| Carbendazim | 105.7 | 10.8 | 100.1 | 10.6 | 90.5 | 7.6 |
| Thiabendazole | 91.2 | 4.9 | 87.9 | 6.8 | 85.0 | 4.0 |
| Pyrimethanil | 112.2 | 2.7 | 107.0 | 3.2 | 102.8 | 4.9 |
| Cyprodinil | 104.3 | 3.2 | 99.9 | 6.1 | 100.2 | 4.9 |
| Diazinon | 104.9 | 5.6 | 102.0 | 6.6 | 99.2 | 6.8 |
| Pyrazophos | 99.9 | 4.0 | 96.6 | 5.6 | 91.3 | 4.1 |
| Chlorpyrifos | 91.7 | 4.6 | 99.5 | 5.2 | 97.2 | 3.8 |

| Pesticide | Sample 1 | Sample 2 | Sample 3 | Sample 4 | Sample 5 | Sample 6 |
|---------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Methamidophos | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |
| Carbendazim | < 1.5 | < 1.5 | < 1.5 | 10.2 | 8.7 | 2.3 |
| Thiabendazole | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |
| Pyrimethanil | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |
| Cyprodinil | 1.7 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |
| Diazinon | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |
| Pyrazophos | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |
| Chlorpyrifos | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 | < 1.5 |

Pesticides detected in red wine samples (ng/mL)